

Table 17.2a. Relative standard errors (%) of total above-ground biomass on forest land by forest type, age class and boreal zone in Canada.

Boreal zone	Forest type	Age (years)											
		1-20	21-40	41-60	61-80	81-100	101-120	121-140	141-160	161-180	181-200	201+	Other ¹
Non-boreal	Treed Mixedwood	5.8	5.4	3.6	4.0	5.3	9.7	12.3	17.3	26.2	59.9	27.3	0.0
	Treed Coniferous	6.8	6.8	5.3	5.3	5.0	5.3	5.8	6.3	6.9	8.3	3.9	0.0
	Treed Broadleaf	6.0	6.8	4.5	4.5	5.1	11.1	14.1	23.7	43.3	45.2	57.6	0.0
	Non-treed	51.6	99.9	86.7	2.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.0
Boreal	Treed Mixedwood	6.2	5.8	4.9	3.5	2.9	5.1	9.2	13.6	31.9	51.5	36.9	0.0
	Treed Coniferous	5.0	4.5	3.8	2.7	1.7	1.9	3.0	6.0	8.8	12.7	8.0	0.0
	Treed Broadleaf	7.9	7.5	5.0	4.1	5.1	8.6	16.9	41.2	39.2	100.0	74.8	0.0
	Non-treed	51.5	49.6	45.8	55.4	40.2	44.8	100.0	0.0	56.6	0.0	0.0	3.1
Canada	Treed Mixedwood	4.4	4.0	3.0	2.6	2.6	4.5	7.4	10.8	22.7	46.1	25.9	0.0
	Treed Coniferous	4.1	3.8	3.1	2.5	1.6	1.8	2.8	4.4	5.4	7.3	3.6	0.0
	Treed Broadleaf	4.8	5.1	3.4	3.1	3.6	6.8	10.9	20.5	30.4	44.1	45.6	0.0
	Non-treed	37.7	49.5	42.7	55.4	40.2	44.8	100.0	0.0	56.6	0.0	0.0	2.7

Source: Canada's National Forest Inventory, **revised 2006 baseline** (Version 3, December 2013) (<http://nfi.nfis.org>)

Note: The Non-boreal zone in the Arctic ecozones (Arctic Cordillera, Northern Arctic, Southern Arctic) are not inventoried. The total land and water area amounts to approximately 252,202.08 thousand ha.

¹ Missing or unknown age class.