

Table 18.0a. Relative standard errors (%) of total above-ground biomass on forest land by species group and age class in Canada.

Species group	Age (years)											
	1-20	21-40	41-60	61-80	81-100	101-120	121-140	141-160	161-180	181-200	201+	Other ¹
Unclassified	13.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.6
Unspecified hardwoods	7.4	17.6	29.3	60.6	99.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other hardwoods	11.3	10.2	10.4	9.5	19.7	23.7	32.7	73.2	64.9	0.0	0.0	0.0
Maple	11.2	8.2	5.1	6.1	7.1	13.9	17.4	27.9	61.0	100.0	85.1	0.0
Birch	12.5	7.0	4.6	4.6	5.6	10.1	18.9	66.4	56.4	90.5	100.0	0.0
Poplar	6.8	5.3	3.8	3.0	3.8	6.6	10.6	21.1	23.8	62.9	36.9	0.0
Unspecified conifers	5.3	8.2	20.5	54.5	73.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cedar & other conifers	39.7	14.4	12.6	9.8	12.1	13.2	25.9	25.4	32.2	34.4	8.5	0.0
Larch	11.1	10.5	11.9	13.1	12.4	15.6	18.8	21.6	25.0	41.9	33.0	0.0
Douglas-fir	39.7	23.1	17.6	14.6	13.0	9.9	12.8	12.2	15.3	20.7	13.2	0.0
Hemlock	43.9	18.0	16.0	18.1	18.1	20.1	26.2	19.7	21.1	21.8	6.1	0.0
Fir	11.6	5.7	4.8	6.0	5.4	7.7	10.0	10.6	10.6	12.3	5.9	0.0
Pine	7.9	6.4	5.6	4.3	4.3	5.6	6.5	8.3	11.4	15.9	12.6	0.0
Spruce	6.4	4.7	3.4	2.6	1.4	1.8	2.8	5.6	7.4	12.0	6.1	0.0

Source: Canada's National Forest Inventory, **revised 2006 baseline** (Version 3, December 2013) (<http://nfi.nfis.org>)

Note: The Arctic ecozones (Arctic Cordillera, Northern Arctic, Southern Arctic), and the Taiga Plains and Hudson Plains ecozones located in Nunavut, are not inventoried. These non-inventoried areas total 262,189.89 thousand ha.

¹ Missing or unknown age class.